arXiv:astro-ph/0509603v1 20 Sep 2005

# Particle acceleration in electron-ion jets

K.-I. Nishikawa*, P. Hardee†, C. B. Hededal**, G. Richardson‡, R. Preece§,
H. Sol¶, G. J. Fishman‖, C. Kouveliotou‖ and Y. Mizuno*

*National Space Science and Technology Center, Huntsville, AL 35805
†Department of Physics and Astronomy, The University of Alabama, Tuscaloosa, AL 35487
**Niels Bohr Institute, Department of Astrophysics, Juliane Maries Vej 30, 2100 København O, Denmark
‡Department of Mechanical and Aerospace Engineering University of Alabama in Huntsville Huntsville, AL 35899
§Department of Physics, University of Alabama in Huntsville, Huntsville, AL 35899 and National Space Science and Technology Center, Huntsville, AL 35805
¶LUTH, Observatore de Paris-Meudon, 5 place Jules Jansen 92195 Meudon Cedex, France
‖NASA-Marshall Space Flight Center,
National Space Science and Technology Center, Huntsville, AL 35805

**Abstract.** Weibel instability created in collisionless shocks is responsible for particle (electron, positron, and ion) acceleration. Using a 3-D relativistic electromagnetic particle (REMP) code, we have investigated particle acceleration associated with a relativistic electron-ion jet fronts propagating into an ambient plasma without initial magnetic fields with a longer simulation system in order to investigate nonlinear stage of the Weibel instability and its acceleration mechanism. The current channels generated by the Weibel instability induce the radial electric fields. The $z$ component of the Poynting vector ($\mathbf{E}\times\mathbf{B}$) become positive in the large region along the jet propagation direction. This leads to the acceleration of jet electrons along the jet. In particular the $E\times B$ drift with the large scale current channel generated by the ion Weibel instability accelerate electrons effectively in both parallel and perpendicular directions.

## SIMULATION RESULTS

Recent PIC simulations using injected relativistic electron-ion jets show that acceleration occurs within the downstream jet, rather than by the scattering of particles back and forth across the shock as in Fermi acceleration [1-5, 8].

In this report we present new simulation results of particle acceleration and magnetic field generation for relativistic jet shocks using 3-D relativistic electromagnetic particle-in-cell (REMP) simulations. Our new simulations with further diagnostics has revealed the mechanism of particle acceleration with the Weibel instability.

In the collisionless shock generated behind the head of the relativistic jet the Weibel instability is excited in the downstream region. The instability generates current filaments elongated along the streaming direction and associated transverse loop magnetic fields. These current channels generate the induced radial electric fields which are mainly perpendicular to the loop transverse magnetic fields. Since the Weibel instability is convective, the the current channels propagate with the jets. The $\mathbf{E}\times\mathbf{B}$ also propagate with the jet. Some particles are trapped and accelerated.

Figure 1 (left) shows the $z$ component of $\mathbf{E}\times\mathbf{B}$ with the $x$ and $y$ components of magnetic field (by the arrows) generated by the ion Weibel instability in the $x-y$ plane at $z/\Delta = 430$ for the electron-ion jet injected into an unmagnetized ambient electron-ion plasma at $t = 59.8/\omega_{\rm pe}$. At the region with large positive $z$ component electrons are accelerated along the jet propagation direction. Consequently, at the front part of jet (one third) electrons are accelerated more than $10\gamma v$ as shown in the left panel [6].



**FIGURE 1.** Left and right panels show the $z$ component of $\mathbf{E}\times\mathbf{B}$ at $z/\Delta = 430$ and velocity distributions of jet electrons (One third of jet electrons in the front region is plotted by red curves and on third of rear part of jet is plotted by blue curves) at $t = 59.8/\omega_{\rm pe}$, respectively. Jet electrons are binned as a function of $\gamma v$, where $\gamma = (1-(v_{\rm x}^2+v_{\rm y}^2+v_{\rm z}^2)/c^2)^{-1/2}$.

The $z$ component of $\mathbf{E}\times\mathbf{B}$ becomes mainly positive on $z-x$ plane (not shown). This component accelerate jet electrons effectively as also shown in Fig. 1a in [2]. Certainly, the negative component decelerate particles. This $E\times B$ drift acceleration leads to sufficient acceleration in parallel and perpendicular directions.

We have also performed simulations with widely distributed Lorentz factor of jet electrons and positrons which are assumed to be created by the photon annihilation [7].

## ACKNOWLEDGMENTS

This research (K.N.) is partially supported by the National Science Foundation awards ATM 9730230, ATM-9870072, ATM-0100997, INT-9981508, and AST-0506719. The simulations have been performed on IBM p690 (Copper) at the National Center for Supercomputing Applications (NCSA) which is supported by the National Science Foundation.

## REFERENCES

1. J. T. Frederiksen, C. B. Hededal, T. Haugbølle and Å. Nordlund, *Astrophys. J.* **608**, L13-L16 (2004)
2. C. B. Hededal, J. T. Frederiksen, T. Haugbølle and Å. Nordlund, *Astrophys. J.* **617** L107-L110 (2004)
3. C. B. Hededal and K.-I. Nishikawa, *Astrophys. J.* **623**L89-L92 (2004)
4. K.-I. Nishikawa, P. Hardee, G. Richardson, R. Preece, H. Sol, and G. J. Fishman, *Astrophys. J.* **595**, 555-563 (2003)
5. K.-I. Nishikawa, P. Hardee, G. Richardson, R. Preece, H. Sol, and G. J. Fishman, *Astrophys. J.* **622**, 927-937 (2005)
6. K.-I. Nishikawa, P. Hardee, C. Hededal and G. J. Fishman, *Astrophys. J.*, in preparation (2005)
7. K.-I. Nishikawa, E. Ramirez-Ruiz, P. Hardee, G. J. Fishman, and C. Kouveliotou, *Astrophys. J.*, in preparation (2005)

8. L. O. Silva, R. A. Fonseca, J. W. Tonge, J,.M. Dawson, W. B. Mori, and M. V. Medvedev, *Astrophys. J.* **596**, L121-L124 (2003)